

Unfavorable Ocean Conditions Likely Cause Of Low 2007 Salmon Returns Along West Coast

Date: March 4, 2008

Source: NOAA National Marine Fisheries Service

Summary: Scientists are reviewing unusual environmental conditions in the Pacific Ocean as the likely culprit for the dramatically low returns of Chinook and coho salmon to rivers and streams along the West Coast of the United States in 2007.

FULL STORY

NOAA scientists are reviewing unusual environmental conditions in the Pacific Ocean as the likely culprit for the dramatically low returns of Chinook and coho salmon to rivers and streams along the West Coast of the United States in 2007.

Researchers from NOAA's Northwest and Southwest Fisheries Science Centers are comparing data on the low food production of the California Current in 2005 that occurred when this year's returning salmon would have been entering the ocean from their natal streams to feed and grow.

The cold waters of the California Current flow southward from the northern Pacific along the West Coast and are associated with upwelling, an ocean condition caused by winds that bring nutrients to the ocean's surface and is the main source of nourishment for the ocean's food web. In 2005 a southward shift in the jet stream, delayed favorable winds and upwelling for the California Current, which normally begins in spring. The winds instead arrived in mid-July, causing high surface water temperatures and very low nutrient production within the near-shore marine ecosystem.

"We are not dismissing other potential causes for this year's low salmon returns," said Usha Varanasi, NOAA Fisheries Service Science Center Director for the Northwest Region. "But the widespread pattern of low returns along the West Coast for two species of salmon indicates an environmental anomaly occurred in the California Current in 2005."

New data released by the Pacific Fisheries Management Council indicate the 2007 returns of fall Chinook salmon to the Sacramento River in California's Central Valley were approximately 33 percent of what fishery biologists expected. Projections for 2008 are substantially lower than last year's estimate.

Coho salmon returning to spawning streams in California and Oregon are also considerably lower than predicted. A preliminary analysis found an average 27 percent of the parental stock returning in 12 streams monitored in California. Even though coho returns appear to improve along the coast from south to north, Oregon Coast coho salmon had less than 30 percent of their parental stock return.

Coho salmon are listed as either endangered or threatened under the Endangered Species Act in the Central/Northern California and Southern Oregon watersheds.

Story Source:

Materials provided by **NOAA National Marine Fisheries Service**. *Note: Content may be edited for style and length.*

Cite This Page:

MLA

APA

Chicago

NOAA National Marine Fisheries Service. "Unfavorable Ocean Conditions Likely Cause Of Low 2007 Salmon Returns Along West Coast." ScienceDaily. ScienceDaily, 4 March 2008.
<www.sciencedaily.com/releases/2008/03/080303145253.htm>.

RELATED STORIES**Resident Orcas' Appetite Likely Reason for Decline of Big Chinook Salmon**

Dec. 16, 2019 — Large, old Chinook salmon have mostly disappeared from the West Coast. A new study points to the recent rise of resident killer whales, and their insatiable appetite for large Chinook salmon, as the ...

West Coast Waters Returning to Normal but Salmon Catches Lagging

Mar. 9, 2018 — Ocean conditions off most of the US West Coast are returning roughly to average, after an extreme marine heat wave from about 2014 to 2016 disrupted the California Current Ecosystem and shifted many ...

Largest Chinook Salmon Disappearing from West Coast

Feb. 27, 2018 — The largest and oldest Chinook salmon -- fish also known as 'kings' and prized for their exceptional size -- have mostly disappeared along the West Coast, according to a new ...

Dramatic Decline in Genetic Diversity of Northwest Salmon

Jan. 10, 2018 — Columbia River Chinook salmon have lost as much as two-thirds of their genetic diversity, researchers have found. The researchers reached this conclusion after extracting DNA from scores of bone ...

Featured Content*from New Scientist***Stunning photograph shows a flower as an insect might see it**

April 13, 2022 — Debora Lombardi's image, shortlisted for the 2022 Sony World Photography Awards, uses a photography technique to show the fluorescence of flowers under UV light, revealing them as insects might see them.

Fabric conditioners reduce the release of microfibres in tumble dryers

April 6, 2022 — Tumble-drying clothes produces microfibre pollution, but this is almost halved by using a tumble dryer sheet and an anti-wrinkle fabric conditioner.

Tropical city air pollution led to 470,000 premature deaths in 2018

April 8, 2022 — Cities in the tropics are experiencing a growing air pollution problem, which is estimated to have led to a 62 per cent rise in premature deaths since 2005.

Visit *New Scientist* for more global science stories >>>